Label-free Profiling of Ligands for Endogenous GPCRs Using a Cell-Based High Throughput Screening Technology

Ye Fang*, Gary Li, and Ann M. Ferrie

Science and Technology Division, Corning Incorporated, *Life Sciences*, Corning, NY 14831
Email: fangy2@corning.com

Epic™ system

CORNING
Discovering Beyond Imagination



Abstract

G protein-coupled receptors (GPCRs) are the largest family of cell surface receptors and the largest class of drug targets. The activation of GPCRs is known to lead to the dynamic translocation of multiple signaling molecules or molecular assemblies during its signaling cycle, and in many cases cytoskeletal reorganization. Such a movement and/or reorganization results in dynamic redistribution of cellular contents, equivalent to dynamic mass redistribution (DMR), which can be monitored online in living cells using Corning® Epic™ system – a label-free and non-invasive biosensor system that is centered around resonant waveguide grating (RWG) biosensors. The resultant DMR signal offers a novel and functional optical signature for studying GPCR signaling and screening GPCR drug compounds. The theoretical considerations and assay principles are first described. The unique optical signatures for over 20 endogenous G_i, G_s and G_q-coupled receptors in A431 human epidermoid carcinoma cells are presented. As exampled in studying the signaling pathways and network interactions of endogenous GPCRs, as well as screening GPCR ligands using both kinetic and endpoint measurements, Corning® Epic™ system is an easily scaleable biosensor HTS platform for G_i, G_s and G_q GPCR drug discovery and deorphanization.

Cell-Based Assays on the Epic™ System: Theory of Operation

- Evanescent waveguide sensor
- Resonant coupling of light into a waveguide using a diffraction grating
- Waveguide formed of four layers, each with different index of refraction
- Monitor changes in reflected wavelength or angle (proportional to local mass redistribution)
- Only monitor the bottom portion of cells:
  - Evanescent wave has limited penetration into cell monolayer on the sensor surface - depth/sensing volume (~120nm)



Cell-Based Assays on the Epic™ System: Dynamic Mass Redistribution (DMR)

The spatial and temporal redistribution in local mass density and amount in the vicinity of the sensor surface

- cellular targets or molecular assemblies moving into or out of the sensing volume of the optical biosensor

"There's Plenty of Room at the Bottom"
-- Richard P. Feynman

Cell-Based Assays on the Epic™ System: Assay Principles



Publications

Fang, Y., Li, G., and Peng, J. (2005) Optical biosensor provides insights for bradykinin B_2 receptor signaling in A431 cells. *FEBS Lett.* 579, 6365-6374.

Fang, Y., Ferrie, A.M., Fontaine, N.H., and Yuen, P.K. (2005) Characteristics of dynamic mass redistribution of EGF receptor signaling in living cells measured with label free optical biosensors. *Anal. Chem.* 77, 5720-5725.

Fang, Y., Ferrie, A.M., and Li, G. (2005) Probing cytoskeleton modulation by optical biosensors. *FEBS Lett.* 579, 4175-4180.

Fang, Y., Ferrie, A.M., Fontaine, N.H., and Yuen, P.K. (2005) Optical biosensors for monitoring dynamic mass redistribution in living cells mediated by epidermal growth factor receptor activation. *Proceedings of IEEE Engineering in Medicine and Biology, 361 (pp1-4).*

Fang, Y., Ferrie, A.M., Li, G., and Fontaine, N.H. (2006) Cellular functions of cholesterol probed with optical biosensors. *In Press.*

Fang, Y., Ferrie, A.M., and Fontaine, N.H. (2006) Resonant waveguide grating biosensors for living cell sensing. Submitted.

Fang, Y., and Ferrie, A.M. (2006) Signaling of protease-activated receptors in A431 cells: dynamic mass redistribution and regulation by cholesterol. *Submitted.*

List And Profiles of Agonists Examined for Known *Endogenous GPCRs* in A431

Receptor	Signaling	Agonist	K_d (Epic™)*	K_d (Flexo-3)	Reference
A1 (ADORA1)	G_i	Adenosine-amine-congener	3.7±1.1μM	N.A.	Klaus, H. E. *et al.* (1997) *J. Biochem. Mol. Biol. Biophys.* 1, 109-116
A2A (ADORA2A), A2B (ADORA2B)	G_s	NECA	72.4±28.6nM	N.A.	Klaus, H. E. *et al.* (1997) *J. Biochem. Mol. Biol. Biophys.* 1, 109-116
β1 (ADRB1)	G_s	Epinephrine	5.8±1.8nM	N.A.	Delorme-Kluchlin, C. *et al.* (1984) *FEBS Lett.* 169, 151-155.
B2 (BDKRB2)	G_q	bradykinin	6.5±1.6nM	10.0±2.8nM	Liebmann, C. *et al.* (1996) *Biochem. J.* 313, 109-118.
BB1 (BRS1)	G_q	Bombesin	N.D.	N.D.	Corbleyn, Y. *et al.* (1994) *FEBS Lett.* 348, 280-284.
H1 (HRH1)	G_q	HTMT	45.4±14.2nM	N.A.	Tilly, B.C. *et al.* (1990) *J. Cell Biol.* 110, 1311-1315.
H2 (HRH2)	G_s				
LPA1	G_i	LPA			Ohta, H. *et al.*(2003) *Mol. Pharmacol.* 64, 994-1005.
LPA2, LPA3	G_q		N.D.	524±164nM	
P2Y1 (hex), P2Y1, P2Y6	G_q	ATP	2.2±0.6μM	7.0±1.7μM	van Rhal, J. *et al.* (2005) *J. Biol. Chem.* 280, 9870-9878.
P2Y2	G_q				Gong, A.Y.M. *et al.* (2003) *J. Immun. Determatol.* 121, 335-327.
P2Y11	G_s G_q				van Rhal, J. *et al.* (2005) *J. Biol. Chem.* 280, 9870-9878.
PAR1	G_q $G_{12/13}$	Thrombin SFFLR-amide	9.6±2.8nM/ml 1.8±0.1μM	6.0±1.0nM/ml 5.0±0.4μM	van Rhal, J. *et al.* (2005) *J. Biol. Chem.* 280, 9870-9878.
PAR2	G_q	Trypsin GLEGR-amide SLIGKV-amide	980±27.8nM 2.3±0.4μM 6.1±1.0μM	45.7±5.8nM 2.5±0.2μM 3.8±0.4μM	Ivankot, R. *et al.* (2004) *J. Invest. Dermatol.* 122, 937-944.

* The K_d value was determined based on the amplitude of P-DMR signal as a function of agonist concentration

List of Agonists Examined for GPCRs That Are Unknown for Their Endogenous Level in A431

Receptor	Signaling	Agonist	Receptor	Signaling	Agonist
M1, M1,M5	G_q	Oxotremorine-M	GAL2	G_q	galanin
M2, M4*	G_i	oxotremorine-M	GAL2	G_q	galanin
α2A (ADRA1A)	G_q	oxymetazoline	GCGR	G_s	glucagon
α2A, B, C (ADRA2A,B,C)	G_i	UK14304, Clonidine	Y1,2,4,5,6 (NPY1,2,4,5,6R)	G_i	neuropeptide Y
β1 (ADRB1)	G_s	Xamterol	MC1,2,3,4,5R	G_s	β-MSH, α-MSH
AT1 (AGTR1)	G_q/G_i	angiotensin II	MOTR	G_i	morfin
BB2,3 (BRS1,2)	G_q	Bombesin	NTSR1	G_q	Neurotensin, neurotensin N
CB1,2 (CNR1,2)	G_i	Anandamide	μ (OPRM1)	G_i/o	dynorphin A, DAMGO
CXCR1	G_i	IL-8	κ(OPRK1)	G_i/o	dynorphin A
CXCR3	G_i	IP-10	ORL1 (OPRL1)	G_i/o	nociceptin
D1 (DRD1)	G_s	A-77636, SKF38393	5-HT$_{1A}$ (HTR1AF)	G_i/o	8BOAH
D5 (DRD5)	G_s	SKF 38393	NK1 (TACR1)	G_q	substance P
ET-A,B (EDNRA,B)	G_q	endothelin-1	NK2 (TACR2)	G_q	neurokinin A
AP1R (AGTR2.1)	?	apelin 1-17	UT2 (GRP14)	G_q	urotensin II
GAL1,3	G_i/o	galanin	V1a, V1b	G_q	vasopressin

* Signatures were obtained with Epic™ system.

Dynamic Mass Redistribution in Cell Biology: Location, Location, Location – Signaling and Trafficking

- Cellular targets are highly organized and spatially restricted
- The relocalization of proteins and molecular assemblies is fundamental to:
  - cell signaling by enabling precise control in its amplitude, duration and kinetics
  - cell functions – migration, invasion, growth, cycling, differentiation, survival, and death

There are many cellular events that lead to mass redistribution

- **Protein target or substrate recruitment**
  - to the nucleus
  - to the membrane
  - to the cytosol
  - throughout recycling pathways
  - to or from other organelles
  - uptake from extracellular space
- **Endocytosis and recycling**
- **Exocytosis**
- **Apoptosis**
- **Cytoskeleton rearrangement**

Huddler, A., *et al.* (2003) *Mol. Cell. Biol.* **23**, 9318-9326
Fang, Y., *et al.* (2005) *FEBS Lett.* **579**, 4175-4180
Kholodenko, B.N. (2003) *J. Expar. Biol.* **206**, 2073-2082.

Signature of G_q-Coupled Receptors: Thrombin Receptor



The Nature of DMR Mediated Through G_q-Coupled Receptors: Thrombin Receptor



Signature of G_s-Coupled Receptors: β2 Receptor



Switching of GPCR Signaling: $LPA_{1,2,3}$ Receptors in A431 Cells



- G_q signaling occurs within the range of LPA doses examined.
- The DMR signals exhibited a switching from primarily G_i at low LPA doses to G_q at high LPA doses.

Ohta, H. *et al.*(2003) *Mol. Pharmacol.* 64, 994-1005.

Mapping Signaling Network Interactions: Bradykinin B_2 Receptor in A431 Cells



Neves et al. *Science* 2002, 296, 1636.
Graness et al. *Biochem. J.* 2000, 347, 441-7
Liebmann et al. *Biochem. J.* 1996, 313, 109-118.

Mapping Signaling Network Interactions: Bradykinin B_2 Receptor in A431 Cells



- B_2 receptor in A431 uses dual signaling pathways: G_q and G_s.
- The G_s and G_q signaling can cross-regulate each other.

Receptor Activation and Desensitization: Protease-Activated Receptors in A431 Cells



Regulation of GPCR Signaling by Cholesterol: Cholesterol Depletion with Methyl-β-cyclodextrin



Compound Potency



Partial Agonist-Mediated GPCR Activation



High-Throughput GPCR Cell-Based Assays: End-Point Measurements

- End-point measurements
- Offline solution addition
- 384well Epic™ plate
- 5 minutes assay



Conclusions

- Epic™ System is also a scalable biosensor HTS platform for ***biochemical assays***.
- Epic™ System is a scalable biosensor HTS platform for ***cell-based G protein-coupled receptors drug discovery and deorphanization***.
- The Epic™ biosensor provides novel and unique signatures for all G_i-, G_s-, and G_q-coupled receptors.
  - Label-free
  - Standard SBS 96 well and 384 well microplate format
  - Endogenous GPCRs
  - Real physiological conditions
  - Multiplexing
  - High information content:
    - kinetics, efficacy/potency, signaling pathways, network interactions, signaling regulation, oligomerization, cross-communications, …